Plenvu Instructions

provided, fill with

water, mix and

2- Then drink 2 full containers of water.

drink.

3 DAYS BEFORE 1 DAY BEFORE **7 DAYS BEFORE 5 DAYS BEFORE PROCEDURE PROCEDURE PROCEDURE PROCEDURE** (Prep Day Part 1) *NO SOLID FOOD Do not take any Do not take any Do not take any medicines listed medicines listed medicines listed **TODAY!** below 5 days before below 3 days before below 7 davs before vour procedure: vour procedure: your procedure: You can have clear liquids all 10:00PM: day today. (See STOP the attached clear liquid diet sheet). The more you Plavix Xarelto Aspirin drink, the better Coumadin Pradaxa Ibuprofen your prep will be. Warfarin Eliquis Motrin Savaysa Aleve 5:00PM: Meloxicam PART 1: Mix and Advil drink the 1st dose Ride home: **A member of our Naproxen of Plenvu following •BC Goody's Confirm you staff will call to the instructions confirm your have a provided by the arrival time prior to responsible office. your procedure. If adult to drive a message is left, Mixing Instructions: you home after please return the 1- Pour Dose 1 in the procedure. the clear call at your earliest container You may **NOT** ride the convenience.*

transit or take a taxi

home.)

1 DAY BEFORE PROCEDURE (Prep Day Part 2)

PART 2: Mix and drink the 2nd dose of Plenvu following the instructions provided by the office.

Mixing Instructions:

- 1- Pour Dose 2 pouch A and Pouch B in the clear container, mix with water and drink.
- 2- Then drink 2 full containers of water.
 - **You may continue clear liquids until the allowed time in the next column.**

PROCEDURE DAY

YOU CAN HAVE CLEAR LIQUIDS TODAY UNTIL:



(No gum, candy, mints, ice chips, sips of clear liquid or chewing tobacco after the time listed above.)

Early morning:

Take your usual prescribed heart medication / blood pressure medication with a sip of water if applicable. Do not take any other medications before your procedure unless instructed to do SO.